



New York State Department of Health

2001-02 EDITION

covering data reported
through December 1999

Additional copies of this report may be obtained by writing to:

New York State Department of Health
Bureau of HIV/AIDS Epidemiology
Empire State Plaza
Albany, NY 12237

Additional and related information is also available from the New York State Department of Health Web site on the Internet:
<http://www.health.state.ny.us>.

Comments regarding the format or content of this report are welcome.

In an effort to reduce the costs of printing, please notify the producer of this document if you wish your name to be deleted from our mailing list or if your address has changed.

AIDS

IN NEW YORK STATE

2001-2002 EDITION

covering data reported
through December 1999

State of New York
Department of Health

Introduction

AIDS in New York State provides a wide range of statistical information on the HIV / AIDS epidemic in New York State based largely on data collected by the New York State Department of Health. It highlights findings from epidemiologic studies, surveillance and program evaluations. As such, it serves as a valuable resource for those concerned with monitoring and addressing the epidemic.

The trend in reduction in the number of deaths from AIDS seen over the past few years in New York and across the nation continues. Many people living with HIV and AIDS continue to benefit from new pharmaceuticals and treatments, living longer, healthier lives. The decrease in HIV infection among women of childbearing age and newborns also continues. This is the good news.

The decline in AIDS cases and deaths can be attributed in part to the wide range of innovative health care and supportive services for people living with HIV, put into place in New York over the past eighteen years. These services were developed through the creative use of state, local, federal and private funding and with a unique collaboration among government, health and community providers, and affected communities.

In 1998, legislation requiring a system of HIV / AIDS surveillance was passed; this system was implemented on June 1, 2000. With the availability of HIV surveillance data, the epidemiological focus will shift from AIDS to HIV, providing a much more current profile of the HIV epidemic in New York State. The next issue of *AIDS in New York State* will highlight these new HIV data. An interim report on HIV surveillance data is anticipated to be published in 2002.

There is still much work to be done. The HIV / AIDS epidemic in New York State continues to disproportionately impact communities of color, particularly African Americans and Latinos. Young adults continue to be diagnosed with AIDS in their twenties, indicating probable infection during their teen years. In some populations, young women have higher infection rates than men.

All those who work in and are affected by HIV / AIDS and public health will want to keep informed about these and other trends in the epidemic. *AIDS in New York State* is a key resource for all people interested in monitoring the epidemic.

Table of Contents

HIV Infection in New York State	1
Section 1	HIV Infection in Childbearing Women 3
Figure 1-1	New York State Survey of Childbearing Women HIV Prevalence by Region 4
Figure 1-2	New York State Survey of Childbearing Women HIV Prevalence by Race/Ethnicity 9
Figure 1-3	New York State Survey of Childbearing Women HIV Prevalence by Age 12
Figure 1-4	New York State Survey of Childbearing Women HIV Prevalence by County of Residence 15
Figure 1-5	New York State Survey of Childbearing Women HIV Prevalence by Zip Code of Residence in New York City 17
Section 2	HIV Infection in Sexually Transmitted Disease (STD) Clinic Clients 18
Figure 2-1	Unlinked Serosurvey of STD Clients HIV Prevalence in Three Continuously Participating Clinics Outside New York City by Risk Behaviors 19
Figure 2-2	Unlinked Serosurvey of STD Client HIV Prevalence in Three Continuously Participating Clinics Outside New York City – All Clients 20
Figure 2-3	Unlinked Serosurvey of STD Clients HIV Prevalence in Three Continuously Participating Clinics Outside New York City – Male Clients 22
Figure 2-4	Unlinked Serosurvey of STD Clients HIV Prevalence in Three Continuously Participating Clinics Outside New York City – Female Clients 24
Figure 2-5	STD Clinic Clients Who Seek HIV Testing HIV Positivity by Public Health Region and Gender 26
Section 3	HIV Infection in Adolescents and Young Adults 28
Figure 3-1	Unlinked Serosurvey of Adolescent Clinic Clients HIV Prevalence in One New York City Clinic 30
Figure 3-2	Unlinked Serosurvey of Homeless Youth Clinic Clients HIV Prevalence in One New York City Clinic 32
Figure 3-3	HIV Prevalence Among Military Recruits by Demographic Characteristics and Region 36

Figure 3-4	HIV Prevalence Among Job Corps Entrants by Demographic Characteristics and Region	40
Figure 3-5	Youth Requesting HIV Testing at the Office of Children and Family Services-Operated Residential Facilities for Court Adjudicated Youth	42
Section 4	HIV Infection in Substance Users	45
Figure 4-1	Unlinked Serosurvey of Methadone Maintenance Treatment Programs HIV Prevalence in Three Clinics in New York City	46
Figure 4-2	Unlinked Serosurvey of Methadone Maintenance Treatment Programs HIV Prevalence in Three Clinics in New York City - Male Clients	48
Figure 4-3	Unlinked Serosurvey of Methadone Maintenance Treatment Programs HIV Prevalence in Three Clinics in New York City - Female Clients	50
Figure 4-4	Unlinked Serosurvey of Methadone Maintenance Treatment Programs HIV Prevalence in Three Clinics in New York City by Borough of Residence	52
Figure 4-5	Unlinked Serosurvey of Drug Treatment Clinic Clients HIV Prevalence in Five Clinics in New York State	56
Figure 4-6	Unlinked Serosurvey of Drug Treatment Clinic Clients HIV Prevalence in Five Clinics in New York State - Male Clients	58
Figure 4-7	Unlinked Serosurvey of Drug Treatment Clinic Clients HIV Prevalence in Five Clinics in New York State - Female Clients	60
Figure 4-8	Drug Treatment Client Voluntary Testing Program HIV Positivity by Client Demographics, IDU Status and Residence of Client	62
Section 5	HIV Infection in Prison Inmates	65
Figure 5-1	State Prison Inmate Voluntary HIV Testing Program HIV Positivity by Client Demographics, IDU Status and Residence of Client	66
Section 6	HIV Infection in Medical Patients	69
Figure 6-1	HIV Status of Tuberculosis Cases by Sex and Region	70
Figure 6-2	HIV Prevalence in Medical Practice Specimens Tested by the New York State Laboratory	71
Figure 6-3	Community Health Clinics HIV Positivity by Client Sex, Age, Race/Ethnicity and Exposure Category of Those Tested	72
Section 7	State-Operated Anonymous Test Sites	77
Figure 7-1	State-Operated Anonymous Test Site Clients HIV Positivity by Sex and Location of Test Site	78
Figure 7-2	State-Operated Anonymous Test Site Clients – All Sites in New York State HIV Positivity by Sex, Race/Ethnicity and Exposure Category	82
Figure 7-3	State-Operated Anonymous Test Site Clients - New York City Sites HIV Positivity by Sex, Race/Ethnicity and Exposure Category	88

Section 8	Planning Resources	89
	Community Need Index	89
Table 8-1	Counties Included in the Eight Regional Reports and the Three CNI statistical areas	89
Table 8-2	Definitions of the Ten Health Statistical Indicators Used in the CNI Composite Measure	91
Figure 8-1	Community Need Index Report Series (2000 Edition) Sample County Table	92
Figure 8-2	Community Need Index Scores and Location of NYSDOH AIDS Institute-Sponsored Providers Sample Map	93
	HIV Counseling and Testing Resource Directory	94
	Behavior Risk Surveys	95
	Youth	95
	Adults	96
	Information on the Web	97
Section 9	AIDS Morbidity and Mortality Overview	99
Figure 9-1	AIDS Incidence by Year of Diagnosis	102
Figure 9-2	States That Have Reported 5,000 or More AIDS Cases	103
Figure 9-3	Cumulative AIDS Cases by Exposure Category, Race/Ethnicity, Gender and Age at Diagnosis	104
Figure 9-4	AIDS Cases Confirmed by Exposure Category, Race/Ethnicity, Gender and Age at Diagnosis	106
Figure 9-5	AIDS Cases and Rates per 100,000 Population Confirmed in 1998 and 1999 and Cumulative Totals by Age Group in the United States, New York State and Surrounding States	108
Figure 9-6	AIDS Cases by Exposure Category and Year of Diagnosis	109
Section 10	Geographic Distribution of AIDS Cases in New York State	111
Figure 10-1	AIDS Cases and Rate per 100,000 Population by County and City of Residence and Year of Diagnosis	112
Figure 10-2	Cumulative AIDS Incidence Rates per 100,000 Population by County and City of Residence	114
Figure 10-3	AIDS Cases by Ryan White Region and Year of Diagnosis, Cumulative Cases and Rate per 100,000 Population	115
Figure 10-4	Adult AIDS Cases by Age at Diagnosis, Gender, Race/Ethnicity, Exposure Category and Ryan White Region	116
Figure 10-5	AIDS Cases by Exposure Category, Year of Diagnosis and Ryan White Region	120
Figure 10-6	AIDS Cases Diagnosed in 1998 and 1999 per 100,000 Population by Race and Ryan White Region	134
Figure 10-7	AIDS Cases Diagnosed 1996-1999 by Exposure Category, Year of Diagnosis and Ryan White Region	137
Figure 10-8	AIDS Cases Diagnosed 1996-1999 in New York State exclusive of New York City by Age at Diagnosis, Gender, Exposure Category, Race/Ethnicity and Region	139
Figure 10-9	AIDS Cases Among New York City Residents by Exposure Category, Race/Ethnicity and Borough	140

Figure 10-10	Adult AIDS Cases by Age at Diagnosis, Gender Race/Ethnicity, Exposure Category and New York City Borough	142
Figure 10-11	Adults Living with AIDS, and AIDS Case Rates by United Hospital Fund Neighborhood System in New York City	144
Figure 10-12	AIDS Cases Diagnosed 1996 – 1999 in New Jersey Counties Abutting New York City by Age at Diagnosis, Race/Ethnicity, Gender and Exposure Category .	145
Section 11	Age at Diagnosis of AIDS	147
Figure 11-1	Cumulative Adult AIDS Cases by Race/Ethnicity, Age at Diagnosis, Gender, Exposure Category and Region	148
Figure 11-2	Adult AIDS Cases by Race/Ethnicity, Age at Diagnosis, Gender, Exposure Category and Region	150
Figure 11-3	Adult AIDS Cases Diagnosed 1996 – 1999 by Age at Diagnosis, Gender, Exposure Category, Race/Ethnicity and Region	154
Section 12	Pediatric AIDS Cases	155
Figure 12-1	Distribution of Pediatric AIDS Cases Diagnosed 1996 - 1999 by Age at Diagnosis, Race/Ethnicity, Gender and Exposure Category	156
Figure 12-2	Pediatric AIDS Cases Diagnosed 1996 – 1999 by Race/Ethnicity, Exposure Category, Age at Diagnosis, Case Rate per 100,000 Population and Residence	157
Figure 12-3	Pediatric AIDS Cases by Year of Diagnosis	158
Figure 12-4	Cumulative AIDS Cases Among Children Infected Perinatally by Race/Ethnicity, Age at Diagnosis, Residence, Current Age and Mother’s Exposure Category	159
Figure 12-5	Selected Diseases at Diagnosis Among Pediatric AIDS Cases by Year of Diagnosis	160
Section 13	AIDS Cases Among New York State Adolescents and Young Adults	161
Figure 13-1	AIDS Cases Among Adolescents and Adults Under Age 25, Diagnosed 1996 – 1999 by Age at Diagnosis, Race/Ethnicity, Gender and Exposure Category	162
Figure 13-2	AIDS Cases Among Adolescents and Adults Under Age 25, Diagnosed 1996 – 1999 by Exposure Category, Year of Diagnosis, Gender, Race/Ethnicity, Age at Diagnosis and Residence	163
Figure 13-3	AIDS Cases Among Adolescents and Adults Under Age 25 1996 – 1999 by Gender, Exposure Category and Race/Ethnicity	164
Figure 13-4	AIDS Incidence Among Adolescents and Adults Under Age 25, Diagnosed by Exposure Category, Race/Ethnicity and Year of Diagnosis	165
Section 14	AIDS Among New York State Adults	166
Figure 14-1	AIDS Prevalence: Number of Adults Alive with AIDS at the End of Each Year, by Gender	167
Figure 14-2	Adults Living with AIDS by Race/Ethnicity, Age at Diagnosis, Gender, Exposure Category and Region	168
Figure 14-3	Adult Female AIDS Cases Diagnosed in 1998 and 1999 by Race/Ethnicity and Exposure Category	170
Figure 14-4	Adult Female AIDS: Cumulative Cases by Year of Diagnosis	171

Figure 14-5	Adult Female AIDS Cases Diagnosed 1996 – 1999 by Race/Ethnicity, Year of Diagnosis, Exposure Category, Age at Diagnosis and Residence	172
Figure 14-6	Adult Male AIDS Cases Diagnosed in 1998 and 1999 by Race/Ethnicity, and Exposure Category	173
Figure 14-7	Adult Male AIDS: Cumulative Cases by Year of Diagnosis	174
Figure 14-8	Adult Male AIDS Cases Diagnosed 1996 – 1999 by Race/Ethnicity, Exposure Category, Year of Diagnosis, Age at Diagnosis and Residence	175
Figure 14-9	AIDS Cases Diagnosed 1996 – 1999 Among People Aged 50 and Older by Exposure Category, Year of Diagnosis, Gender, Race/Ethnicity, Age at Diagnosis and Residence	176
Section 15	HIV Transmission Categories and Subpopulations	177
Figure 15-1	Distribution of AIDS Cases by Age at Diagnosis, Race/Ethnicity, Gender and Exposure Category	180
Figure 15-2	AIDS Incidence Among Men Who Have Sex With Men by Race/Ethnicity, Year of Diagnosis and Region	182
Figure 15-3	AIDS Cases Among Men Who Have Sex With Men by Race/Ethnicity, Year of Diagnosis and Region	183
Figure 15-4	AIDS Incidence Among Injecting Drug Users by Race/Ethnicity, Year of Diagnosis and Region	184
Figure 15-5	AIDS Incidence Among Injecting Drug Users by Race/Ethnicity, Year of Diagnosis and Region	185
Figure 15-6	AIDS Cases Attributable to Heterosexual Transmission by Race/Ethnicity, Year of Diagnosis, Gender, Age at Diagnosis, Exposure Category of Partner and Residence	186
Figure 15-7	AIDS Cases Attributable to Heterosexual Transmission by Race/Ethnicity and Year of Diagnosis	187
Section 16	AIDS Cases Among Inmates in New York State Correctional Facilities	189
Figure 16-1	Distribution of AIDS Cases Among State Prison Inmates	190
Figure 16-2	AIDS Cases Diagnosed 1996 – 1999 Among State Prison Inmates By Race/Ethnicity, Year of Diagnosis, Gender, Age at Diagnosis and Exposure Category	192
Section 17	Mortality Due to AIDS	193
Figure 17-1	Mortality Among AIDS Cases by Year of Diagnosis	194
Figure 17-2	Mortality Among AIDS Cases by Year of Diagnosis	195
Figure 17-3	Deaths from HIV / AIDS in New York City	196
Figure 17-4	Ranking of AIDS Among the Five Leading Causes of Death by Population Group and Age	197
Figure 17-5	AIDS as a Leading Cause of Death for New York City Men and Women Ages 20-39	198
Section 18	Health Services and Expenditures for HIV/AIDS in New York State HIV Uninsured Care Programs	199
Figure 18-1	ADAP Enrollment by Year of Enrollment	201
Figure 18-2	ADAP Enrollment by Gender, Race/Ethnicity and Year of Enrollment	202

Figure 18-3	ADAP Drug Utilization by Category-Users by Month	203
Figure 18-4	ADAP Antiretroviral Utilization	204
Figure 18-5	ADAP Combination Antiretroviral Utilization by Race/Ethnicity (Three+ ARV)	205
Figure 18-6	ADAP Combination Antiretroviral Utilization by Risk (Three+ ARV)	205
Figure 18-7	ADAP Combination Antiretroviral Utilization by Gender (Three + ARV)	206
Figure 18-8	HIV Uninsured Care Programs Actual Expenditures	207
Section 19	Use of Hospitals for HIV/AIDS Care	208
Figure 19-1	HIV/AIDS Hospital Discharges, Days and Length of Stay by Year	210
Figure 19-2	HIV/AIDS Average Daily Inpatient Census	212
Figure 19-3	Pediatric HIV/AIDS Hospital Discharges, Days and Length of Stay by Year and Region	214
Figure 19-4	HIV/AIDS Tuberculosis Hospitalizations	216
Figure 19-5	HIV/AIDS Inpatient Statistics by Hospital Location	218
Figure 19-6	HIV/AIDS Inpatient Statistics by Hospital Location	220
Figure 19-7	HIV/AIDS Hospital Discharges by Disposition	221
Figure 19-8	HIV/AIDS Inpatient Statistics by Hospital Auspice	222
Figure 19-9	HIV/AIDS Inpatient Statistics for Designated AIDS Centers, HHC Hospitals and Other Hospitals	224
Figure 19-10	HIV/AIDS Hospitalizations by Primary Expected Payor by Year	226
Section 20	HIV/AIDS-Related Costs and Expenditures in New York State	229
Figure 20-1	Estimated Total Expenditures for Persons with HIV/AIDS by Payor	230
Figure 20-2	Estimated Annual Expenditures for Persons with HIV/AIDS	232
Figure 20-3	Estimated HIV/AIDS Funding by State Agency and Program New York State Funds	233
Figure 20-4	Annual Medicaid Expenditures for People with HIV/AIDS	234
Figure 20-5	Medicaid Expenditures for People with AIDS	236

HIV INFECTION IN NEW YORK STATE

Due to the extended incubation period between HIV infection and clinical illness, AIDS cases can reflect HIV transmission patterns that occurred a decade or more ago. To gain more complete and timely data on the prevalence of HIV infection among New York's citizens, the New York State Department of Health monitors HIV prevalence in several ways:

Unlinked HIV Seroprevalence Studies

Unlinked HIV seroprevalence studies test leftover blood specimens, originally drawn for routine diagnostic tests, for HIV antibodies, after all other testing has been completed and all personal identifying information has been removed from the specimen. The seroprevalence studies are performed exclusively for epidemiologic purposes and the test results cannot be linked back to the individual from whom the specimen was originally drawn. This type of study reduces the selection bias that occurs when people volunteer to be tested. Data on demographic characteristics, clinical conditions and risk behaviors are obtained from the client's charts prior to the blinding and testing of the specimen. However, since clients are not interviewed, data on risk behavior is often very limited. A wide variety of sites have been studied using this methodology, including sexually transmitted disease (STD) clinics, substance abuse treatment clinics, adolescent clinics, homeless

youth clinics and prison intake centers.

Much of this section of AIDS in New York has been modified from previous editions to provide more information on trends in HIV prevalence during the last several years. Keep in mind when reviewing the following charts and graphs, that while most of the charts indicate a decline in **prevalence** over the years, this does **not** necessarily mean that the rate of new infections (**incidence**) has also dropped. For example, the rate of new infections in a population may remain stable, while the number of infected people sampled in a study may decline. An example of this occurring is when a new facility, devoted to persons with specific risk activities, opens and draws clients away from a survey clinic, thus lowering the prevalence seen in the survey clinic. Conversely, an increase in prevalence does not necessarily indicate an increase in the number of new infections. With more HIV infected individuals receiving treatment and living longer, prevalent cases may be encountered during the course of a study, without any new infections having occurred.

Data from unlinked HIV seroprevalence studies conducted by the New York City Department of Health, Office of AIDS Research is available directly from that office. The publication *HIV Seroprevalence Update* is available on the web at <http://www.ci.nyc.ny.us/html/doh/home.html>, or by contacting

the New York City Department of Health, Bureau of Disease Intervention Services, Office of AIDS Research/HIV Serosurveys, 346 Broadway, New York, NY 10013.

Voluntary and Anonymous Counseling and Testing Data

The Department of Health monitors the results of people voluntarily seeking HIV testing through confidential and anonymous counseling and testing sites, medical clinics, private physicians' offices, prisons and drug treatment programs.

National Seroprevalence Studies that include New York State Residents

The Department of Defense and the Job Corps test new applicants for HIV and have made their prevalence data available without identifying information.

Caveats

There are several caveats that should be considered when interpreting the HIV prevalence data:

- Sampling bias cannot be eliminated. For example, the HIV prevalence among females attending an upstate STD clinic cannot be generalized to all females in areas outside of New York City, as those who attended the clinic may be more likely to be at risk for HIV infection. Even prevalence studies of

entire populations must be interpreted with care. The results from the Survey of Childbearing Women, for example, cannot be generalized to other reproductive-age women, because they represent only women carrying pregnancies to term. Similarly, the results from the military survey cannot be generalized to all young adults because those engaging in certain risk behaviors may not apply for military service and, therefore, may “self-select” out of the sample.

- In certain instances the number of individuals with specific characteristics may be too small to permit meaningful interpretations, or to maintain confidentiality of those tested. In these instances, groups or categories may be combined to create new categories, often termed “other/unknown”. A specific example of this is with regards to Asian/Pacific Islanders and

Native American/Alaskan Natives. When possible, data for these groups have been provided. However, in some prevalence studies, the small number of individuals tested has prohibited release of this information.

- When reading and interpreting tables it is important to remember that results obtained from larger studies can be used with more confidence than those generated from a study with a smaller number of participants. To highlight study sizes, HIV prevalence data in this year’s publication is provided in tables that correspond to the graphs. Please be sure to read the tables that accompany the graph. Note the study size on each table (number tested) and the scale on the x-axis of each graph. Remember to generalize only to the population from which the sample was drawn.

Prevalence of HIV infection within the groups studied in New York State varied significantly in 1999, from 23 percent among clients of New York City methadone maintenance treatment programs to 0.14 percent among women giving birth in areas outside New York City.

Overall, these data demonstrate a high correlation between injecting drug use and HIV infection in both men and women. Prevalence of HIV infection is also elevated among men and women who are being treated for other STDs. In general, the prevalence of HIV infection is consistently higher among residents of New York City compared with the rest of the state, and is highest in people between the ages of 20 and 49, and among blacks and Hispanics.